

ITP-SB-99-36
hep-th/9907142
July, 1999

Fundamental vs. Solitonic Description of D3 branes

I.Y. Park ¹,

C.N. Yang Institute for Theoretical Physics
State University of New York
Stony Brook, N. Y. 11794-3840

Abstract

Type IIB string theory expanded around D3 brane backgrounds describes the dynamics of D3 branes as solitonic objects. On the other hand, there is a fundamental description of them via Polchinski's open strings with Dirichlet boundary conditions. Since these two descriptions describe the dynamics of the same objects, D3 branes, it is natural to believe that they are dual. Therefore at this level, we have a string-string duality as opposed to a string-field theory duality. Once we take the same limits in both descriptions, Maldacena Conjecture in its weaker form follows. We try to make this viewpoint precise and study the implication of it for the stronger form of Maldacena Conjecture.

¹email: ipark@insti.physics.sunysb.edu

In the progress of string theory during the past several years, D-branes [1] have played a central role. In particular, the authors of [2, 3, 4, 5] studied the absorption of scalars by D3 branes and noticed that supergravity calculations of the cross sections for various scattering processes are reproduced by the gauge theories on the branes. More recently, by studying the extremal 3 brane configurations of IIB supergravity Maldacena made a conjecture [7]: The weaker form of the conjecture states that IIB supergravity on $\text{AdS}_5 \times \text{S}_5$ is holographically [8, 9] dual to large N , $\mathcal{N}=4$, $d=4$ Super Yang-Mills (SYM_4) theory and the stronger form states that IIB *string* theory on $\text{AdS}_5 \times \text{S}_5$ is dual to SYM_4 for any value of g_s and N . There are several reviews of the topic, for example, [10, 11, 12, 13].

Large amount of literature has appeared so far to support the weaker form of the conjecture. However, not much has been done along the line of proving or deriving the conjecture. In this article, we address the question of why the weaker form of the conjecture should be true. Similar reasoning appeared in [14]. However the viewpoint we take does not lead to the stronger form. Instead we seem to obtain a string-string duality. More specifically, we propose and try to make precise the following: IIB string theory on $\text{AdS}_5 \times \text{S}_5$ should be dual to oriented open string theory¹ propagating in the same background.

A couple of years ago, it was shown [15] that closed strings (or supergravity in the low energy) describe the dynamics of D-branes at length scales much larger than the string scale while it is the open strings (or world volume theory in the low energy) which dominate at the sub-stringy scales. Then one may ask whether one can put together open strings, closed strings and branes in one theory such that the theory is valid for the entire range of length scale.

This is a very natural question because D-branes are, in some sense, topological defects that convert closed strings into open strings when the close strings touch the branes [2, 3, 4, 5, 6]. Then there should be reversed processes in which open strings become closed strings and leave the branes. We take these properties of D-branes as our axiomatic starting point: The “physical” system under consideration consists of D-branes, open strings and closed strings, and we look for theories that describe their dynamics. Below we argue that there are (at least) two ways to combine these three

¹The closed strings produced by open string interactions are understood throughout the article.

objects (although there are some subtleties) and they correspond to two different stringy descriptions of D3 brane² dynamics: one is “fundamental” and the other is “solitonic”. In the solitonic approach, one looks for a theory in which the D-branes are embedded as solitons while one would consider the coordinates of the branes as dynamical fields in the fundamental description. For convenience and reasons which will become more clear, we will use the same word “embedding” for both descriptions.

It is well known that type IIB supergravity has extremal solutions of parallel 3-brane configurations [17, 18]. For simplicity we only consider N coincident D-branes in this article. The solutions preserve one half of the supersymmetry and carry Ramond-Ramond (RR) charges. These facts allow one to view these particular solutions as a low energy realization of the D3 branes introduced by Polchinski. Even with the full string corrections, one expects to have corresponding solutions: a D3 brane will be a soliton of the full IIB string theory. Type IIB string theory expanded around them provides a solitonic description of D3 brane dynamics, Fig 1(a). In this description, D3 branes appear as solutions of field equations. They interact with each other by exchanging the perturbative spectrum of IIB theory excited “around” them. The situation is very similar to that of monopoles in ordinary field theories, where one considers perturbative excitations around a monopole solution.

Then the subtlety lies in the way the open strings appear in this description. In other words, if IIB string theory expanded around a D3 brane solution correctly describe the dynamics (we assume it does), how does it incorporate the open strings? To answer this question precisely will require much work. However, given the fact that the open strings are associated with the excitations of the branes, a natural guess will be that the combined system of branes and open strings on them should be somehow associated with solutions with various non-extremality. We will return to this matter later. In summary, we started with IIB closed strings, the D branes appear as solitons, open string effects will be embedded in some solutions with energies higher than those of extremal solutions and IIB string theory expanded around the D3 brane solutions describes their dynamics as solitons.

Now we seek another way of putting together the open strings, closed strings and D3 branes. Since we started with closed strings above, we start with D3 branes this time. One might consider the supermembrane Lagrangian [19] introducing the

²In this article, we only consider D3 branes which are non-singular [16]. For D_p branes with $p \neq 3$, there may be subtleties with the singularities.




Figure 1: (a) D3 branes appear as backgrounds in the solitonic description. (b) D3 branes appear as Dirichlet boundary conditions (represented by the dotted lines) in the fundamental description.

coordinates for the branes and try to quantize the theory. However, a direct quantization of such extended objects is known to be a difficult task [20, 21]. Quantization of bosonic membranes has been discussed more recently in [22]. Fortunately one can achieve the same goal by an alternative description through open strings to which we now turn.

Obviously the open strings must be oriented and have Dirichlet boundary conditions since the open strings we need are exactly the same open strings produced when the closed strings touch the D-branes. In this description, D branes are introduced as boundary conditions, Figure 1(b). (D branes are not merely boundary conditions, but they should provide a background in which open strings propagate. We will discuss this point later.) Finally closed strings are produced through the interactions of open strings, which will describe the reversed processes mentioned previously.

It has been argued by Witten [23] that in the low energy, the dynamics of D3 branes are described by SYM_4 theory. SYM_4 has scalar fields which have the interpretation of the transverse coordinates³ of the D-branes. Therefore the open string description can be considered as a fundamental description of D branes. It is a *fundamental* description since the coordinates of the branes appear in the Lagrangian as

³There may be higher derivative corrections to the identification of the scalars and the transverse coordinates of the branes [24].

dynamical variables.

Above we have found two different ways of combining the three objects: open strings, closed strings and D3 branes. These in turn correspond to two different descriptions of the same D3 brane system: one is a solitonic description of IIB string theory for the D3 brane configuration under consideration, and the other is an open string description with Dirichlet boundary conditions. This is exactly the concept of duality: the two descriptions must be dual since they describe the same D3 brane system, which must have unique physics, at the same energy scale, the string scale. Therefore before taking any limits, we have a string-string duality. We will call this “fundamental-solitonic duality”. In the solitonic description, closed strings are “primary” objects while the open strings are associated with some excited states of 3-brane configurations. On the other hand, it is the open strings which are primary in the fundamental description and the closed strings appear through the interactions of the open strings. Below we take this viewpoint seriously and study the implications of it.

The organization of this paper is as follows: First, we briefly demonstrate that the weaker form of Maldacena conjecture follows from the fundamental-solitonic duality by taking the same limits on both sides of the two descriptions. Then we argue that in high energy, one should expect a string-string duality rather than the stronger form of the Maldacena conjecture which is a string-field theory duality. These two points can be made without finding the precise statement of the fundamental-solitonic duality. Then we try to make this string-string duality more precise and study the additional implications of it. Finally we conclude with a summary and open problems.

To show that our viewpoint leads to the weaker form of Maldacena Conjecture, we consider N coincident D3 branes, Figure 2. Once again, it should be emphasized that we take limits *starting from the string-string duality*. Other than the fact that we do not take any limit for “ r ”, the coordinate transverse to the branes⁴, the argument should be rather familiar and we will be brief. Figure 2(b) represents the fundamental description and Figure 2(a) the solitonic one. Consider the low energy limit where one can neglect the massive string modes: the effective actions of both sides contain only *massless* string modes. We also impose the following common limits⁵ on both sides, $N \rightarrow \infty, g_s \rightarrow 0, g_s N = \text{fixed but very large}$. First consider the description (b), the

⁴We will discuss the near-horizon limit of r later.

⁵Similar limits were taken in [3].




Figure 2: We take common limits in Solitonic Description, (a) and Fundamental Description, (b)

fundamental description. According to Witten [23], the low energy effective action of N coincident D3 branes is $U(N)$ gauge theory. Even though $g_s \rightarrow 0$, we can not neglect the quantum corrections (because the effective loop expansion parameter is $g_s N$), but rather the theory becomes strongly coupled. Therefore the resulting theory in the description (b) is $\mathcal{N}=4$, $d=4$ $U(N)$ SYM with quantum correction.

This is in contrast with the solitonic description where the theory becomes classical. The metric configuration for N coincident D3 branes is [17]

$$\begin{aligned}
 ds^2 &= H^{-\frac{1}{2}} dx_4^2 + H^{\frac{1}{2}} (dr^2 + r^2 d\Omega_5^2) \\
 H &= 1 + \frac{4\pi g_s N \alpha'^2}{r^4}
 \end{aligned} \tag{1}$$

In the limit where $g_s N$ is very large, the geometry of the solitonic description becomes $AdS_5 \times S_5$. The radius of curvature of S_5 , R , is given by $R^2 = \alpha' \sqrt{4\pi g_s N}$. The massive stringy corrections are also neglected since they are expected to come as a geometrical series of $\frac{\alpha'}{R^2} = \frac{1}{\sqrt{4\pi g_s N}}$ which becomes negligibly small in the limit we are taking. Therefore we have classical supergravity on $AdS_5 \times S_5$.

Since we started with two dual descriptions, the limiting theories must be dual too: type IIB supergravity on $AdS_5 \times S_5$ is holographically dual to large N strongly coupled $SU(N)^6$ SYM₄ and the precise form of the connection between the two theories was

⁶It has been argued [26, 27] that the bulk theory corresponds to $SU(N)$ part of the $U(N)$ gauge group.

proposed in [25, 26].

Now we turn to the implication of our viewpoint for the stronger form of the Conjecture. It is necessary to keep only the massless modes of open string description to obtain the SYM. Since we must impose the same limit for the solitonic description, it seems to be impossible to have full IIB *string* theory with all the massive modes if our viewpoint is correct. In other words, without taking the limits discussed above one will have oriented open *string* theory as a theory dual to IIB string theory. Therefore, instead of the stronger Conjecture, we propose below to consider oriented open string theory⁷ as a theory dual to IIB string theory on $\text{AdS}_5 \times \text{S}_5$.

Having argued that the fundamental-solitonic duality implies $\text{AdS}_5/\text{CFT}_4$ duality in its weaker form, we now turn to the more precise statement of this string-string duality itself and study other implications of it. For this, we consider Fig 3. From now on, we do not take any limits of g_s, α' and N unless otherwise specified.



Figure 3: (a) Solitonic Description. (b) Fundamental Description

The solitonic description, i.e., type IIB string theory on a background of N coincident D3 branes with RR charge [30], is complicated at the technical level, given the fact that even the string theory on a simpler background such as $\text{AdS}_5 \times \text{S}_5$ is not completely understood although there has been some progress [31, 32, 33, 34]. The D3 branes appear as backgrounds interacting each other by exchanging IIB strings, Figure 3 (a). It is not very clear how this description incorporates the open strings precisely. In [35, 36], calculations of entropy of D-brane system using statistical mechanics of massless open string gas agreed with the Bekenstein-Hawking entropy of

⁷Extension of the weaker form of Maldacena Conjecture to a duality between supergravity and DBI action was discussed in [28, 29].

near-extremal solutions of supergravity. Therefore a natural guess would be that the massive open strings should be associated with non-extremal solutions with large non-externality. In other words, when closed strings are incident on D-branes, open strings appear on the branes, but in the IIB description the combined system of the branes and the open strings on them may appear as some excited states and whether the open strings are massless or massive will be associated with the size of the non-extremality parameter, typically called r_o . Reversely, a brane in its excited state will decay emitting a closed string.

To go to the fundamental description, we turn to our axiomatic starting point: When a closed string touches a D brane, it becomes an open string. This is the open string we must consider to obtain the fundamental description. In this description, the branes are embedded as Dirichlet boundary conditions, Fig 3(b). Since the branes are oriented, the open strings must be oriented⁸. Open string interactions will produce closed strings which must be IIB closed strings. The end points of open strings are constrained to move in a flat hyper-plane⁹. However, the “body” of them propagates in the background produced by D branes. Therefore the D branes are not merely the boundary conditions but they also embed themselves as a background geometry for the open strings to propagate in.

We have just argued that IIB theory on D3 brane backgrounds must be dual to open string theory on the same background. Now we return to the near horizon limit $r \ll R$. This limit allows one to drop 1 in H of eq (1) (without taking any limits for g_s and N), and the metric becomes that of $AdS_5 \times S_5$ again. Therefore we have IIB string theory on $AdS_5 \times S_5$ for the solitonic description. Then we have to consider the same background for the open string description: we have open strings propagating in $AdS_5 \times S_5$. Therefore it is natural to believe that IIB string theory on $AdS_5 \times S_5$ with N units of flux of 5-five form field strength should be dual to an open string theory in the same background¹⁰. Then one immediate question is whether the open

⁸There has been some recent discussion of oriented open and closed string theory in the frame work of string field theory [37].

⁹In the low energy limit where we have a very poor resolution, the open strings on the flat hyper-plane will appear as *particles* moving in four dimensional Minkowski space (the flat hyper-plane) and this is the space where SYM theory is defined.

¹⁰ It has been argued [38, 39] that the Dirac-Born-Infeld plus Wess-Zumino action should be expanded around AdS_5 rather than a flat background to obtain the dimension six operator predicted by supergravity [40, 41, 38]. This is natural according to our viewpoint.

string description has the same symmetries as the IIB description. At the level of DBI action plus WZ term, superconformal symmetry has been discussed in [42]. We expect to have superconformal symmetry at the level of full string theory too. After all, the symmetry should be viewed as coming from the background in the open string description too as it is true in the solitonic description.

Conclusion

In this article, the “physical” system of N coincident D3 branes, open strings and closed strings was considered. We argued that there are (at least) two ways to embed the system in string theory. The two ways correspond to two different descriptions of D3 branes: fundamental and solitonic. Since they are the descriptions of the same system, it is natural to believe that the two descriptions are dual. Taking appropriate, common limits on both sides of the descriptions leads to the weaker form of Maldacena Conjecture. However, we argued that in high energy, it is more natural to expect a string-string duality than the stronger form of Maldacena conjecture which takes the form of a string-field theory duality. In particular, we propose that IIB string theory on $\text{AdS}_5 \times S_5$ with N units of the flux of five-form field strength should be dual to oriented open string theory in the same background.

There are many details to fill in. Many of them are related to the poor status of string theory in various curved backgrounds. In particular, it will be very helpful to learn about oriented open string theory in the geometry of D3 branes or $\text{AdS}_5 \times S_5$. It will also be interesting to study how precisely IIB description incorporates open string effects. We hope to make progress along these lines in near future.

Acknowledgments

I thank J. Maldacena, H. Oogury and J. Schwarz for conversations and S.R. Das, S. de Alwis, J. Polchinski and B. Zwiebach for communications. I am very grateful to I.R. Klebanov for reading the draft and making useful suggestions and corrections. I thank P. van Nieuwenhuizen for raising interesting questions. Finally I would like to thank I. Chepelev, B. Kulik, M. Roček and specially F. Gonzalez-Rey for very helpful discussions and corrections.

References

- [1] J. Polchinski, “TASI Lectures on D-branes”, hep-th/9611050; String Theory Vol. 1,2, Cambridge
- [2] A. Hashimoto and I.R. Klebanov, “Decay of Excited D-branes”, hep-th/9604065
- [3] I.R. Klebanov, “World Volume Approach to Absorption by Non-dilatonic Branes”, hep-th/9702076
- [4] S.S. Gubser and I.R. Klebanov and A.A. Tseytlin, “String Theory and Classical Absorption by Threebranes”, hep-th/9703040
- [5] S.S. Gubser and I.R. Klebanov, “Absorption by Branes and Schwinger terms in the World Volume Theory”, hep-th/9708005
- [6] I.R. Klebanov, “D-branes and Creation of Strings”, hep-th/9709160
- [7] J. Maldacena, “The Large N Limit of Superconformal Field Theories and Supergravity”, hep-th/9711200
- [8] G. 't Hooft, “Dimensional Reduction in Quantum Gravity”, gr-qc/9310026
- [9] L. Susskind, “The World as a Hologram”, hep-th/9409089
- [10] I. Klebanov, “From Threebranes to Large N gauge Theories”, hep-th/9901018
- [11] M.R. Douglas and S. Randjbar-Daemi, “Two Lectures on AdS/CFT Correspondence”, hep-th/9902022
- [12] J.L. Petersen, “Introduction to the Maldacena Conjecture on AdS/CFT”, hep-th/9902131
- [13] O. Aharony, S.S. Gubser, J. Maldacena, H. Oogury, and Y. Oz, “Large N Field Theories, String Theory and Gravity”, hep-th/9905111
- [14] I.R. Klebanov, Unpublished
- [15] M.R. Douglas, D.Kabat, P. Pouliot and S.H. Shenker, “D-branes and Short Distances in String Theory”, hep-th/9608024

- [16] G.W. Gibbons and P.K. Townsend, “Vacuum Interpolation in Supergravity via Super p-branes”, hep-th/9307049
- [17] G. Horowitz and A. Strominger, Nucl. Phys. B360 (1991)197
- [18] M. Duff, R.R. Khuri and J.X. Lu, “String Solitons”, Phys. Rep.259(1995)213, hep-th/9412184
- [19] E. Bergshoeff, E. Sezgin and P.K. Townsend, Phys. Lett. 189B (1987) 75; Ann. Phys. 185 (1988) 330
- [20] B. de Wit, M. Lüscher and H. Nicolai, Nucl. Phys. B320 (1989)135
- [21] B. de Wit, “Supermembranes and Super Matrix Models”, hep-th/9902051
- [22] M. Kaku, “Ultra-Violet Behavior of Bosonic Quantum Membranes”, hep-th/9907016
- [23] E. Witten, “Bound States of Strings and p-branes”, hep-th/9510135
- [24] F. Gonzalez-rey, B. Kulik, I.Y. Park and M. Roček, “Selfdual Effective Action of N=4 Super Yang-Mills”, hep-th/9810152
- [25] S.S. Gubser, I.R. Klebanov and A.M. Polyakov, “Gauge Theory Correlators from Non-critical String Theory”, hep-th/9802109
- [26] E. Witten, “Anti-Desitter Space and Holography”, hep-th/9802150
- [27] O. Aharony and E. Witten, “Anti-de Sitter Space and the Center of the Gauge Group”, hep-th/9807205
- [28] S.S. Gubser, A. Hashimoto, I.R. Klebanov and M. Krasnitz, “Scalar Absorption and The Breaking of The World Volume Conformal Invariance”, hep-th/9803023
- [29] S.P. de Alwis, “Supergravity, the DBI Action and Black Hole Physics”, hep-th/9804019
- [30] D. Berenstein and R.G. Leigh, “Superstring Perturbation Theory and Ramond-Ramond Backgrounds”, hep-th/9904104

- [31] I. Pesando, “A κ Gauge Fixed Type IIB Superstring Action on $\text{AdS}_5 \times \text{S}^5$ ”, hep-th/9808020
- [32] R. Kallosh and J. Rahmfeld, “The GS String on $\text{AdS}_5 \times \text{S}^5$ ”, hep-th/9808038
- [33] R. Kallosh and A.A. Tseytlin, “Simplifying Action on $\text{AdS}_5 \times \text{S}^5$ ”
- [34] D. Polyakov, “On the NSR formulation of String Theory on $\text{AdS}_5 \times \text{S}^5$ ”, hep-th/9812044
- [35] S.R. Das and A.D. Mathur, “Excitations of D-strings, Entropy and Duality”, hep-th/960115
- [36] S.S. Gubser, I.R. Klebanov and A.W. Peet, “Entropy and Temperature of Black 3-Branes”, hep-th/9602135
- [37] B. Zwiebach, “Oriented Open-Closed String Theory Revisited”, hep-th/9705241
- [38] S.R. Das and S.P. Trivedi, “Three Brane Action and The Correspondence Between $\mathcal{N}=4$ Yang Mills Theory and Anti De Sitter Space”, hep-th/9804149
- [39] S. Ferrara, M.A. Lledo and A. Zaffaroni, “Born-Infeld corrections to D3-Brane Action in $\text{ADS}(5) \times \text{S}(5)$ and $\text{N}=4$, $\text{D}=4$ Primary Superfields”, hep-th/9805082
- [40] H.J. Kim, L.J. Romans and P. van Nieuwenhuizen, “The Mass Spectrum of Chiral $\text{N}=2$ $\text{D}=10$ Supergravity on S^5 ”, Phys. Rev. D32 (1985) 389.
- [41] A. Rajaraman, “Two-Form Fields and the Gauge Theory Description of Black Holes”, hep-th/9803082
- [42] P. Claus, R. Kallosh and A. Van Proeyen, “Conformal Symmetry on World Volumes of branes”, hep-th/9812066